

BEST COPY AVAILABLE

Dr. John C. Bugher, M.D., Director
Division of Biology and Medicine

May 13, 1954

Gordon M. Dunning, Health Physicist
Biophysics Branch, Division of Biology and Medicine

401809

ESTIMATED RADIATION DOSE TO THYROID OF NATIVES FROM RONGELAP

This memorandum is in reply to your request for an estimate of additional doses to the thyroid of the Rongelap natives due to the fact that tellurium, as a precursor to iodine, may be present in the gut after ingestion of fallout material. The tellurium, in turn, might disintegrate into radioactive iodine while in the gut, with subsequent deposition of the iodine in the thyroid.

There are some 17 radioactive isotopes of tellurium but only 7 of these are produced in fission. Of these, 6 are not of interest (4 have too short a half-life, 1 leads to stable iodine-127 and 1 leads to iodine-129 with a half-life of 2.4×10^7 years). The remaining radioisotope is tellurium-132, with a half-life of 77 hours leading to iodine-132 with a half-life of 2.4 hours. (Incidentally there is no tellium precursor that is of interest here.)

Without having the original data of LASL I have accepted their estimate that there were ingested and/or inhaled the products of 5×10^{13} fissions, assumed they were all ingested, and then proceeded to calculate the dose to the thyroid from (a) I^{131} (b) each short-lived iodine isotope of interest and (c) the added dose coming from $T^{132}-I^{132}$. The calculations show that $T^{132}-I^{132}$ will produce an added dosage of about 26%.

The best estimated percentage absorption and deposition of iodine is yet to be determined. The best estimate I can turn up to date is still the 20% quoted in NBS Handbook 52. However, I will continue to search for additional information. In the meantime the table below indicates the magnitude of doses to the thyroid if one assumes 20, 50, and 100% absorption and deposition. Incidentally, it may be noted that the calculations below based on 20% (the number assumed by LASL) show estimated doses to the thyroid from I^{131} and from shorter lived iodine isotopes to be in good agreement with those estimated by LASL, i.e., 50 reps for I^{131} and 80 reps for short-lived iodine isotopes.

DEPARTMENT OF ENERGY DECLASSIFICATION REVIEW
SINGLE REVIEW AUTHORIZED BY: DETERMINATION (CIRCLE NUMBER(S))
1 CLASSIFICATION RETAINED
2 CLASSIFICATION CHANGED TO
3 CHANGING NO. DSE CLASSIFIED INFO
4 CO-ORDINATE WITH
5 CLASSIFICATION CANCELLED
6 CLASSIFIED INTO BRACKETED
AA Simigelli - 3/3/74
REVIEWER (ADD):
NAME: K2d
DATE: 4/4/94

This document contains information which is exempt from release under the provisions of the Freedom of Information Act, 5 U.S.C. 552, and is to be controlled in accordance with the provisions of the Atomic Energy Act, 42 U.S.C. 2014, and the Atomic Energy Regulations, 41 CFR 101-11.6.

Access # 326-78
Box 3
File = NMB-B

OFFICE ► BMBP
SURNAME ► DUNNING:MMack. *GMS*
DATE ► 5-13-54

MILITARY RESEARCH & APPL

May 13, 1954

DOSE TO THYROID (REPS)

	Assuming 100% Retention		Assuming 50% Retention		Assuming 20% Retention	
I ¹³¹	255	255	128	128	51	51
I ¹³²	27*	(54)**	14*	(27)**	5*	(10)**
I ¹³³	370	370	185	185	74	74
I ¹³⁵	60	60	30	30	12	12
I ¹³² (Te ¹³²)	185*	(370)**	93*	(185)**	37*	(74)**
Total	897*	(1109)**	450*	(555)**	189*	(221)**

* If assume that one-half of the I¹³² (half-life 2.4 hours) present in the gut is deposited in the thyroid.

** If assume all of the I¹³² (half-life 2.4 hours) present in the gut is deposited in thyroid.

Most probable estimate of ratio of doses to thyroid is:

$$\frac{I^{132}, I^{133}, I^{135}}{I^{131}} \approx 2.5$$

~~THIS DOCUMENT IS UNCLASSIFIED~~

OFFICE ▶ _____

SURNAME ▶ _____

DATE ▶ _____

May 13, 1954

ANNEX

Calculations of Dose to Thyroid

 I^{131}

Assume: inhalation and/or ingestion of 5×10^{13} fissions

At D / 1 there are 0.017 d/m/10 000 fissions

or 8.5×10^7 d/m/ 5×10^{13} fissions

or 38.3 μ c intake of I^{131}

or 1.4×10^{12} atoms intake of I^{131}

(Average energy 0.22 Mev)

$$\text{Dose (reps)} = \frac{(1.35 \times 10^{12}) * (0.22)(1.6 \times 10^{-6})}{(20)(93)} = 255 \text{ reps}$$

* Correction for biological decay.

 I^{132}

Assume: inhalation and/or ingestion of 5×10^{13} fissions

At D / 1, I^{132} intake is 1.1×10^{11} atoms

The average mean energy of I^{132} is about 0.55 mev or 2.5 times that of I^{131} .

Thus, the energy equivalent to I^{131} would be

$$(1.1 \times 10^{11})(2.5) = 2.75 \times 10^{11} \text{ atoms of } I^{131}$$

However, due to the short half-life of I^{132} (2.4 hrs) assume that the energy equivalent of 1.5×10^{11} atoms of I^{131} reaches the thyroid.

Thus, the ratio of doses $\frac{I^{131}}{I^{132}} \approx 9.0$

CONFIDENTIAL

OFFICE ► _____

SURNAME ► _____

DATE ► _____

Form AEC-818

U. S. GOVERNMENT PRINTING OFFICE 16-63761-2

501209r

Dr. Bugher

- 4 -

May 13, 1954

I^{133}

Assume: inhalation and/or ingestion of 5×10^{13} fissions

At $D \neq 1$, I^{133} intake is 1.24×10^{12} atoms

The average mean energy of I^{133} is about 0.36 or 163 times that of I^{131} .

Thus, the ratio of doses $\frac{I^{131}}{I^{133}} \approx 0.7$

I^{135}

Assume: inhalation and/or ingestion of 5×10^{13} fissions

At $D \neq 1$, I^{135} intake is 2.36×10^{11} atoms.

The average mean energy of I^{135} is about 0.3 mev or 1.36 that of I^{131} .

Thus, the energy equivalent to I^{131} would be

$(2.36 \times 10^{11})(1.36) = 3.2 \times 10^{11}$ atoms of I^{131} energy equivalent.

Thus, ratio of doses $\frac{I^{131}}{I^{135}} \approx 4.2$

OFFICE

5012081

~~CONFIDENTIAL~~

Dr. Bugher

- 5 -

May 13, 1954



Assume: inhalation and/or ingestion of 5×10^{13} fissions.

At B # 1, Te^{132} intake 100 μc

or 1.5×10^{12} atoms

Assume: the time spent in the gut is 77 hrs.

Then, 7.3×10^{11} atoms of Te^{132} will have disintegrated into 7.3×10^{11} atoms of I^{132} .

The average mean energy of I^{132} is about 0.55 Mev or 2.5 times that of I^{131} .

Thus, the energy equivalent to I^{131} would be

$$(7.3 \times 10^{11})(2.5) = 1.8 \times 10^{12} \text{ atoms of } \text{I}^{131}.$$

However, due to the short half-life of I^{132} (2.4 hrs), assume that only the energy equivalent of 1×10^{12} atoms of I^{131} reaches the thyroid.

Thus, the ratio of doses $\frac{\text{I}^{131}}{\text{I}^{132}(\text{Te}^{132})} \approx 1.4.$

DISTRIBUTION: copy 1A - Dr. Bugher
2A - Dr. Fleming
3A - Biophysics Branch
4,5,6A - B&N Files

~~This material contains information affecting the national defense of the United States within the meaning of the espionage laws, and its transmission or revelation in any manner to an unauthorized person is prohibited by law.~~

OFFICE \rightarrow

SURNAME \rightarrow

5012082